



INVESTOR IN PEOPLE

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

REC'D 17 APR 2003


WIPO PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.




Signed

R. Mahoney

Dated

18 MAR 2003



20 FEB 2002 17279-002000
P01/7700 01 00-0203912.1

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road
Newport
South Wales
NP10 8QQ

1. Your reference

GWS/PJG/24287

2. Patent application number

(The Patent Office will fill in this part)

0203912.1

19 FEB 2002

3. Full name, address and postcode of the or of each applicant (underline all surnames)

Resolution Chemicals Limited
Wedgwood Way
Stevenage
Hertfordshire
SG1 4QT

Patents ADP number (if you know it) 4165429001

If the applicant is a corporate body, give the country/state of its incorporation

4. Title of the invention

Solvent-based sterilisation of pharmaceuticals

5. Name of your agent (if you have one)

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

MATHYS & SQUIRE
100 Gray's Inn Road
London WC1X 8AL
United Kingdom

Patents ADP number (if you know it)

1081001

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number
(if you know it)

Date of filing
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

- a) any applicant named in part 3 is not an inventor, or
 - b) there is an inventor who is not named as an applicant, or
 - c) any named applicant is a corporate body.
- See note (d))

YES

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description

7

Claim(s)

4

Abstract

1

Drawing(s)

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (*Patents Form 7/77*)

Request for preliminary examination and search (*Patents Form 9/77*)

1

Request for substantive examination (*Patents Form 10/77*)

Any other documents
(*please specify*)

-

11.

H. Adams

HARVEY v J ADAMS
MATHYS & SQUIRE

I/We request the grant of a patent on the basis of this application.

Signature

Date

MATHYS & SQUIRE

19 February 2002

12. Name and daytime telephone number of person to contact in the United Kingdom

George W Schlich - 020 7830 0000

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505.*
- Write your answers in capital letters using black ink or you may type them.*
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.*
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.*
- Once you have filled in the form you must remember to sign and date it.*
- For details of the fee and ways to pay please contact the Patent Office.*

SOLVENT-BASED STERILISATION OF PHARMACEUTICALS

5 The present invention concerns a method for the sterilisation of drugs, in particular suspensions of drugs intended for use in nebulizers.

10 Previously it was acceptable for drugs intended for use in nebulizers to be prepared under "clean" conditions. Recently, however, the US FDA has implemented a requirement for all nebulizer solutions to be sterile.

15 In the light of the US FDA decision it is necessary to produce sterile suspension drugs in the US. This is emphasised by problems which have resulted from the use of "clean" suspensions. Multidose formulations made under "clean" conditions were previously acceptable in the US. However such formulations have caused fatalities in the US due to contamination.

20 Drugs typically provided as nebule suspensions are the steroids Fluticasone and Budesonide, used to treat asthma and chronic obstructive pulmonary disorder. These drugs are very insoluble in water and are sold as non-sterile powders.

25 A method of sterilising dry, powdered budesonide is known from International Publication Number WO 99/25359. The method of sterilisation requires budesonide powder to be sterilised and then be mixed with the other components of the formulation under sterile conditions. The drug formulation is subsequently made up under sterile conditions. This method does not permit the complete formulation to be sterilised immediately prior to dispensing into the final sterile vessel.

30 The sterilisation of suspensions raises particular problems. The desired biological activity of the formulation commonly requires that the diameter of particles of the drug lies within a narrow range (typically less than 5 micrometres). The standard means of sterilisation, that is the raising of the temperature of the formulation to 121 °C for 15 minutes, frequently destroys one or more of the components of the formulation. In addition this treatment results in the clumping or agglomeration of the drug particles in the

35

suspension such that the efficacy of the resulting product is impaired or abolished.

5 Known alternative methods for the sterilisation of pharmaceuticals are inappropriate for sterilizing suspension formulations of drugs. Pharmaceuticals may be sterilised by passage through a filter having a pore size of not more than $0.2\mu\text{m}$. However this cannot be used in the case of many suspensions as the required particle size in these formulations is significantly greater than this filter pore size. Similarly, pharmaceuticals may
10 generally be sterilised by gamma-irradiation, but budesonide, for example, is destroyed by such treatment. No further methods for the sterilisation of pharmaceuticals are currently acceptable to regulatory agencies.

15 An object of the present invention is to provide an alternative and/or an improved method for sterilization of suspensions of pharmaceuticals.

20 Accordingly, the present invention provides a method for preparing a sterile composition of a pharmaceutical compound comprising adding solvent to a non-sterile pharmaceutical compound to yield a sterile pharmaceutical compound, optionally removing all or part of the solvent, and under sterile conditions combining the compound with a pharmaceutically acceptable carrier.

25 Hence, in use of the invention sterilization may be achieved by the addition of solvent to the previously non-sterile material. Optional other processing steps may be added, such as filtration as described in more detail below.

30 By sterile, it is meant that resultant pharmaceutical composition meets the requirements of sterility set by medicine regulatory authorities, such as the MCA in the UK or the FDA in the US, and typically the final bio-burden of such compositions is less than 10 CFU/100ml.

35 In a particular embodiment the non-sterile pharmaceutical compound is a powder, for instance a micronised powder. This powder can be mixed with a solvent, forming a combination in which the effect of the solvent is to sterilize the active agent. A steroid in powdered form may be sterilized in this way and particular examples include budesonide or fluticasone.

Generally, any solvent with the required properties may be used in the invention either singly or in combination. It is particularly preferred that the solvent comprises an alcohol. Most particularly preferred is ethyl alcohol, since this solvent is already approved for pharmaceutical formulations for delivery to the lungs. Other alcohols, such as methyl alcohol and isopropyl alcohol, and non-alcohol solvents such as ethyl acetate and TBME may also be used. In order to avoid drug degradation due to hydrolysis of ketal groups, absolute concentrations of solvent (containing no or a minimal water present) are preferred.

Good results have been obtained by adding solvent to the compound at an elevated temperature, suitably from 20°C below the boiling point of the solvent up to its boiling point. This can increase the amount of compound that dissolves and may decrease the dissolving time. Solvent can be added at reflux.

In one aspect of the invention sufficient solvent is used to obtain a slurry of the compound. Hence, there is insufficient solvent to form a solution of the compound, nevertheless sterilization is achieved. An advantage is that only a small quantity of solvent is used and where a minimal amount of solvent is tolerated in the final composition, the solvent added for the purpose of sterilization need not be removed to meet the criteria for acceptance of the final composition for pharmaceutical use.

Where solvent is removed from the sterilized composition, solvent is typically removed under reduced pressure, preferably under vacuum conditions, and preferably at a temperature of 40°C or less. Following evaporation, typically only a few ppm of solvent will remain. If the solvent used is ethyl alcohol, it may be an option to omit this evaporation step, although this is dependent on the final concentration present and maximum allowed levels set by the regulatory bodies. If a non-approved solvent is used, it may be essential to remove all solvent from the final composition.

In a particular embodiment, removing solvent yields a solvent-free, sterile powder, suitable for further processing in a sterile line or for packaging and disposal in this sterile form.

In an alternative aspect of the invention, sufficient solvent is used to dissolve the compound, thus obtaining a solution of the compound. For example, 15 or more volumes of 96% ethanol may be used to dissolve budesonide and 60 or more volumes to dissolve fluticasone.

5

10

15

Optionally, the solution is filtered. The purpose of filtration is as a further sterilisation step and to remove any biomass. Removal of non-viable biomass from the final formulation may not be essential - many pharmaceutical formulations undergo terminal sterilisation, without further filtering. If, therefore, the required level of sterility can be obtained simply by addition of solvent, the filtration step may be unnecessary. Omitting the filtration step would also avoid the expense of replacing clogged filter membranes for every batch of drug solution prepared. In cases where addition of solvent results in incomplete sterilization the addition of a filtration step has the benefit of completing the sterilization

It is an option for the filter, if present, to be sterile. A filter having a pore size of $0.2\mu\text{m}$ or less is preferred.

20

In a yet further aspect of the invention, the sterile pharmaceutical compound is combined with water to form a suspension. If a sterile solution of the pharmaceutical compound is combined with water, the particles will precipitate out of solution to form a suspension. Alternatively, a sterile powder can be added directly to water.

25

The water will typically contain surfactant, such as polysorbate containing compounds, especially Tweens 20 and 80.

30

It is also an option to remove solvent from the suspension. This solvent may be residual from a sterile powder or may be present within the solution before it is combined with water. Solvent is typically removed from the suspension under reduced pressure, preferably under vacuum conditions and preferably at a temperature of 40°C or less.

35

In order to be effective in the lungs, the particle size of the active ingredient, Fluticasone and Budesonide in specific embodiments of the invention, must be within a certain size range, and is likely to need to be reduced following

preparation of a sterile composition as described. Hence, a still further aspect of the invention comprises treating the suspension to obtain a particle size distribution having a mass mean diameter in the range 1-5 μ m, most preferably 2-3 μ m. The sterile suspension may, for instance, be passed through a microfluidiser to reduce the average mass mean diameter of the particles. The microfluidiser is preferably pre-sterilised to avoid contamination of the sterile suspension. A suitable device, referred to as a Microfluidizer^(®), is available from Microfluidics, Inc., described in WO 99/07466.

Further treatments of the sterile suspension can include diluting the suspension with sterile water and/or adding excipients such as EDTA and sodium citrate, so as to obtain the end-product pharmaceutical formulation. Excipients can also be added at an earlier stage, e.g. prior to evaporating solvent or at the same time as adding surfactant.

The sterile composition is then typically stored in ampoules. In use of the invention, the ampoules are nebulisers. Preferably, the nebulisers are pre-sterilised and of the blow-fill-seal type.

The invention further provides a method for preparing a sterile composition of a pharmaceutical compound comprising adding solvent to a suspension of a non-sterile pharmaceutical compound to yield a sterile suspension of a pharmaceutical compound, removing solvent, and under sterile conditions combining the sterile suspension with a pharmaceutically acceptable carrier.

The volume of solvent required to sterilise an aqueous suspension of drug will generally be greater than for sterilisation of a dry powder, therefore thorough evaporation from the suspension is required. Other preferred features of this method are as for the above-described methods of the invention.

The invention also provides a sterile composition of a pharmaceutical compound prepared by adding solvent to a non-sterile pharmaceutical compound to yield a sterile pharmaceutical compound, optionally removing all or part of the solvent, and under sterile conditions combining the compound with a pharmaceutically acceptable carrier.

A sterile composition may be a suspension or a powder.

5 The invention further provides an apparatus for preparing a sterile composition of a pharmaceutical compound, comprising a container defining a sterile inner compartment, a first vessel containing a solvent, and a second vessel containing a non-sterile pharmaceutical compound, arranged so that the solvent can be combined with the non-sterile compound to yield a sterile compound within the compartment, the compartment also containing a sterile aqueous solution into which the sterile compound can be introduced to form a sterile suspension, optionally an apparatus for alteration of the particle size distribution of the suspension and further optionally a sterile exit line for transfer of sterile suspension to sterile ampoules

10 The apparatus may further comprise a sterile filter. Good results have been obtained when the sterile filter of the apparatus has a pore size of 0-2 μ m or less.

15 The invention is now illustrated in specific embodiments by way of the following example.

20 Example 1

Sterilisation of budesonide

25 Powdered Budesonide was dissolved in 15 volumes of 96% alcohol, at reflux. The resultant hot solution was poured through a sterile Pall filter of pore size 0.2 μ m to remove any biomass. 5 litres of sterile water/ Tween 80 concentrate were added per litre of filtered drug solution, causing drug particles to precipitate out of solution, forming a suspension.

30 Evaporation of solvent from the suspension was then carried out at 40°C, under vacuum conditions. An equal volume of water was removed to ensure residual alcohol was kept to a minimum. Residual alcohol concentration was measured and the drug particle size distribution determined.

35 The concentrated suspension was passed through a microfluidiser, using a 100 μ m head, in order to reduce the average drug particle size to a mass

mean diameter of $2-3\mu\text{m}$. Heads of gauge $110\mu\text{m}$, $87\mu\text{m}$ and $75\mu\text{m}$ "Y" and/or "Z" type may also be used, depending on the desired final particle size. The microfluidised suspension was diluted with sterile water - 1 litre suspension being made up to 500 litres - and combined with EDTA and sodium citrate, as excipients for the final formulation. Standard, pre-sterilised nebulisers were filled with diluted drug suspension and sealed ready for use.

Claims

- 5 1. A method for preparing a sterile composition of a pharmaceutical compound comprising adding solvent to a non-sterile pharmaceutical compound to yield a sterile pharmaceutical compound, optionally removing all or part of the solvent, and under sterile conditions combining the compound with a pharmaceutically acceptable carrier.
- 10 2. A method according to Claim 1 wherein the non-sterile pharmaceutical compound is a powder.
3. A method according to Claim 2 wherein the powder is a micronised powder.
- 15 4. A method according to any of Claims 1 to 3 wherein the compound is a steroid.
5. A method according to Claim 4 wherein the compound is budesonide or fluticasone.
- 20 6. A method according to any of Claims 1 to 5 wherein the solvent comprises an alcohol.
- 25 7. A method according to any of Claims 1 to 6, comprising adding solvent to the compound at an elevated temperature, suitably from 20°C below the boiling point of the solvent up to its boiling point.
8. A method according to Claim 7 wherein the solvent is added at reflux.
- 30 9. A method according to any of Claims 1 to 7, comprising using sufficient solvent to obtain a slurry of the compound.
10. A method according to any of Claims 1 to 9, comprising removing solvent under reduced pressure.
- 35 11. A method according to any of Claims 1-10, comprising removing

solvent to yield a solvent-free powder.

5

12. A method according to any of Claims 1 to 11, comprising using sufficient solvent to obtain a solution of the compound.

13. A method according to Claim 12 comprising filtering the solution.

14. A method according to Claim 13 using a filter having a pore size of $0.2\mu\text{m}$ or less.

10

15. A method according to any of Claims 1 to 14, comprising combining the sterile pharmaceutical compound with water to form a suspension.

15

16. A method according to Claim 15, wherein the water contains surfactant.

17. A method according to Claim 15 or 16, comprising removing solvent from the suspension.

20

18. A method according to Claim 17, wherein solvent is removed under reduced pressure.

25

19. A method according to any of Claims 15 to 18, comprising treating the suspension to obtain a particle size distribution having a mass mean diameter in the range $1\text{-}5\mu\text{m}$.

20. A method according to Claim 19, comprising storing the sterile composition in ampoules.

30

21. A method for preparing a sterile composition of a pharmaceutical compound comprising adding solvent to a suspension of a non-sterile pharmaceutical compound to yield a sterile suspension of a pharmaceutical compound, removing solvent, and under sterile conditions combining the sterile suspension with a pharmaceutically acceptable carrier.

35

22. A method according to Claim 21, wherein the compound is a steroid.

23. A method according to any of Claims 21 or 22, wherein the solvent comprises an alcohol.
- 5 24. A method according to any of Claims 21 to 23, comprising adding solvent to the compound at an elevated temperature, suitably not less than 20°C below its boiling point.
- 10 25. A method according to any of Claims 21 to 24, comprising removing solvent under reduced pressure.
26. A method according to any of Claims 21 to 25, wherein the suspension contains a surfactant.
- 15 27. A method according to any of Claims 21 to 26, comprising treating the suspension to obtain a particle size distribution having a mass mean diameter in the range 1-5 μ m.
- 20 28. A method according to Claim 27, comprising storing the sterile suspension in ampoules.
29. A sterile composition prepared according to the method of any of Claims 1-28.
- 25 30. A sterile composition according to Claim 29 wherein the composition is a suspension.
31. A sterile powder prepared according to any of Claims 1 to 11.
- 30 32. Apparatus for preparing a sterile composition of a pharmaceutical compound, comprising a container defining a sterile inner compartment, a first vessel containing a solvent, and a second vessel containing a non-sterile pharmaceutical compound, arranged so that the solvent can be combined with the non-sterile compound to yield a sterile compound within the compartment, the compartment also containing a sterile aqueous solution into which the sterile compound can be introduced to form a sterile suspension, optionally an apparatus
- 35

for alteration of the particle size distribution of the suspension and further optionally a sterile exit line for transfer of sterile suspension to sterile ampoules

- 5 33. Apparatus according to Claim 32, further comprising a sterile filter.
34. Apparatus according to Claim 33, wherein the sterile filter has a pore size of 0-2 μ m or less.

10

15

20

25

30

35

SOLVENT-BASED STERILISATION OF PHARMACEUTICALS

5

ABSTRACT

10

A method for preparing a sterile composition of a pharmaceutical compound comprising adding solvent to a non-sterile pharmaceutical compound and filtering to yield a sterile pharmaceutical compound, optionally removing all or part of the solvent, and under sterile conditions combining the compound with a pharmaceutically acceptable carrier.

15